

2'-Deoxyadenosine-d2

<http://hk.lumiprobe.com/p/2-deoxyadenosine-d2>

2'-Deoxyadenosine-(ribose-5,5-d2) is a deuterated form of the nucleoside 2'-deoxyadenosine and is intended for use as an internal standard for the quantification of deoxyadenosine by LC-MS.

Deoxyadenosine is a nucleoside that is converted to deoxyinosine by the enzyme adenosine deaminase (ADA). Accumulation of deoxyadenosine in the absence of ADA exhibits higher toxicity in lymphoid cells and can inhibit the growth of kinase-deficient cells by inactivating S-adenosylhomocysteine hydrolase (AdoHcyase) and disrupting methylation processes. The triphosphorylated forms of deoxyadenosine analogs are potent inhibitors of ribonucleotide reductase (RNR).



外观:

分子量: 253.26

分子式: $C_{10}H_{11}D_2N_5O_3$

溶解度:

质量控制: NMR 1H 和 HPLC-MS (95+%、D: 98+%)

储存条件:

法律声明: 本產品僅供研究目的提供和銷售。本產品並未經過食品、藥品、醫療器械、化妝品等領域的安全性和效力測試，且未經明示或暗示授權用於其他任何用途，包括但不限於體外診斷、人類或動物用途，以及商業用途。