

DAF-FM (4-Amino-5-Methylamino-2',7'-Difluorofluorescein)

<http://hk.lumiprobe.com/p/diaminofluorescein-daf-fm>

DAF-FM (4-Amino-5-Methylamino-2',7'-Difluorofluorescein) is a cell-impermeant, fluorescent probe for detecting and quantifying low concentrations of nitric oxide (NO). DAF-FM does not need to be activated by cytosolic enzymes and is suitable to detect NO in extracellular matrix.

The fluorescence quantum yield of DAF-FM is ~0.005, but it increases about 160-fold to ~0.81 after reacting with NO and forming a fluorescent benzotriazole (excitation/emission maxima at 495/515 nm) .

The NO detection limit of DAF-FM (~3 nM) is more sensitive than that of DAF-2 (~5 nM). The fluorescence of the NO adduct of DAF-FM is independent of pH above pH 5.5. Moreover, the NO adduct of DAF-FM demonstrates a significantly enhanced photostability compared to that of DAF-2, ensuring reliable results and additional time for imaging.

DAF-FM should be dissolved in DMSO and then used to prepare a working solution. Buffers containing bovine serum albumin (BSA) or phenol red can affect the fluorescence and should be used cautiously.

The cell-permeant version of DAF FM — [DAF-FM DA](#) is also available.

外观: 黄色固体

分子量: 412.35

CAS 编号: 254109-20-1

分子式: $C_{21}H_{14}F_2N_2O_5$

溶解度: 适合甲醇、DMSO、DMF 和水; 水有限; 乙腈和二氯甲烷含量低

质量控制: NMR 1H 和 HPLC-MS (90+%)

储存条件: 收到后 -20°C 避光保存 24 个月。运输: 室温最多可保存3週。干燥。

法律声明: 本产品仅供研究目的提供和销售。本产品并未经过食品、药品、医疗器械、化妆品等领域的安全性和效力测试, 且未经明示或暗示授权用于其他任何用途, 包括但不限于体外诊断、人类或动物用途, 以及商业用途。