


## exo-BCN CE-phosphoramidite

<http://hk.lumiprobe.com/p/exo-bcn-ce-amidite>

BCN-containing phosphoramidite is added at the 5'-terminus of an oligonucleotide. BCN is reactive both to azides (strain-promoted azide-alkyne cycloaddition, SPAAC) and [tetrazines](#) (inverse electron demand Diels-Alder reaction, IEDDA).

Coupling time is standard, like for amidites of natural nucleosides. Exclude the dimethoxytrityl (DMT) removal step and use the Dmt-ON protocol after amidite coupling and oxidation.

Use standard conditions for deprotection and ammonia solution, or AMA mixture (ammonium hydroxide / 40% methylamine, 1:1) for 2 hours at room temperature.



外观: 淡黄色油

质谱 M+ 增量: 343.11

分子量: 481.57

分子式: C<sub>24</sub>H<sub>40</sub>N<sub>3</sub>O<sub>5</sub>P

溶解度: 乙腈良好

质量控制: NMR <sup>1</sup>H、NMR <sup>31</sup>P (95%)

储存条件: 收到後 -20°C 避光保存 12 個月。運輸: 室溫最多可保存3週。乾燥。

法律声明: 本產品僅供研究目的提供和銷售。本產品並未經過食品、藥品、醫療器械、化妝品等領域的安全性和效力測試, 且未經明示或暗示授權用於其他任何用途, 包括但不限於體外診斷、人類或動物用途, 以及商業用途。