

Fluorica-8 AM, green fluorescent calcium indicator

<http://hk.lumiprobe.com/p/fluo-8-am>



Fluorica-8 AM is a cell-permeable Ca^{2+} -indicator that is metabolized by intracellular esterase, leading to a bright green fluorescent signal upon Ca^{2+} -binding (excitation/emission λ at 490/514 nm). Fluorica-8 AM is used for visualization and quantifying intracellular Ca^{2+} . It is well suited for fluorometric and imaging applications such as microscopy, flow cytometry, spectrofluorometry, and fluorometric high-throughput microplate screening assays.

Fluorica-8 AM is similar in structure and spectral properties to the Ca^{2+} indicators Fluo-3 AM and Fluo-4 AM but has the brightest fluorescence compared to them (two times brighter than Fluo-4 and four times brighter than Fluo-3). The K_d of Fluorica-8 AM for Ca^{2+} is about 389 nM. With its highest fluorescence intensity, Fluorica-8 AM is ideal for applications where the concentration of dye loaded into cells is required to be minimized. Unlike Fluo-3 AM and Fluo-4 AM, which require cells to be incubated at 37 °C, Fluorica-8 AM can be loaded into cells at room temperature.

As Fluorica-8 AM does not covalently bind to cellular components, it may be actively effluxed from the cell by organic anion transporters. *In vivo* cell imaging with Fluorica-8 AM is usually performed within one or two hours after loading, but the dye can be re-loaded to cells if it is needed. Fluorica-8 AM can also be fixed *in situ* by [EDC/EDAC](#) for downstream immunofluorescence studies.

Fluorica-8 AM has low solubility in the water. It is recommended to prepare 1 mM stock solution in [labeling grade DMSO](#) prior to cell loading. Use the final concentration of 1-5 μM and incubation at RT for 15-60 min as a start point of your protocol.

Fluorica-8 AM solutions in DMSO may be stored at -20 °C for up to 2 months.



外观:

分子量: 1060.97

CAS 编号: 1345980-40-6

分子式: $\text{C}_{51}\text{H}_{52}\text{N}_2\text{O}_{23}$

溶解度:

质量控制:

储存条件:

法律声明:

本產品僅供研究目的提供和銷售。本產品並未經過食品、藥品、醫療器械、化妝品等領域的安全性和效力測試，且未經明示或暗示授權用於其他任何用途，包括但不限於體外診斷、人類或動物用途，以及商業用途。

激发/吸收极大值, 纳米: 494

