

## Glycine-d2


<http://hk.lumiprobe.com/p/glycine-d2-4896-75-7>

Glycine-d2 is the deuterium-labeled glycine and is intended for use as an internal standard for LC-MS.

Glycine concentration is a good marker of nonketotic hyperglycinemia (genetic, inborn error of glycine metabolism). Nonketotic hyperglycinemia leads to glycine accumulation throughout the body due to a mutation in its cleavage enzyme system. A high level of glycine in the blood may also indicate propionic acidemia or methylmalonic acidemia.

Lower circulating glycine levels could be a marker of metabolic disorders associated with obesity, type 2 diabetes (T2DM), and non-alcoholic fatty liver disease (NAFLDs).

Glycine quantification by LC-MS could be performed most efficiently using glycine-d2 standard.



外观:

分子量: 77.08

CAS 编号: 4896-75-7

分子式: C<sub>2</sub>H<sub>3</sub>D<sub>2</sub>NO<sub>2</sub>

溶解度:

质量控制: NMR <sup>1</sup>H和HPLC-MS (95+ &nbsp;%, d: 98+ &nbsp;%)

储存条件:

法律声明: 本產品僅供研究目的提供和銷售。本產品並未經過食品、藥品、醫療器械、化妝品等領域的安全性和效力測試，且未經明示或暗示授權用於其他任何用途，包括但不限於體外診斷、人類或動物用途，以及商業用途。