


## HMRhoNox-M, Fe(II)-selective fluorescent probe

<http://hk.lumiprobe.com/p/hmrhonox-m>



HMRhoNox-M (also known as LysoRhoNox) is a Fe<sup>2+</sup>-selective fluorescent probe based on the N-oxide-controlled spirocyclization of tetramethyl-hydroxymethyl rhodamine.

In the absence of Fe<sup>2+</sup>, HMRhoNox-M exists in the non-fluorescent spirocyclic form showing only negligible fluorescence in an aqueous buffer and at physiological pH. The addition of Fe<sup>2+</sup> induces a 60-fold increase of the fluorescence signal at 575 nm through the deoxygenation of the dialkylamino group and the transition of the probe to an open fluorescent form. HMRhoNox-M responds to Fe<sup>2+</sup> in a dose-dependent manner.



The fluorescence response of HMRhoNox-M is highly selective for Fe<sup>2+</sup> over other transition metal ions, including Fe<sup>3+</sup>, alkali metal ions, and alkaline earth metal ions.

HMRhoNox-M is the cell-permeant probe that is mainly localized in lysosomes. It is suitable for monitoring fluctuations of endogenous labile iron in living cells, including the transferrin-induced Fe uptake.



外观:

分子量: 388.47

分子式: C<sub>24</sub>H<sub>24</sub>N<sub>2</sub>O<sub>3</sub>

溶解度:

质量控制:

储存条件:

法律声明:

本產品僅供研究目的提供和銷售。本產品並未經過食品、藥品、醫療器械、化妝品等領域的安全性和效力測試，且未經明示或暗示授權用於其他任何用途，包括但不限於體外診斷、人類或動物用途，以及商業用途。

激发/吸收极大值, 纳米: 555

