

(C18) Octadecanoylcarnitine


<http://hk.lumiprobe.com/p/octadecanoylcarnitine>

Octadecanoyl-L-carnitine, also known as stearyl-L-carnitine is a naturally occurring long-chain acylcarnitine. The substance may be used as an analytical reference standard for MS-MS and chromatography-based techniques.

Octadecanoylcarnitine acts as a carrier of long-chain acyl groups from activated fatty acids across the inner mitochondrial membrane, for their further oxidation in the mitochondria. Octadecanoylcarnitine inhibits sodium-dependent reabsorption of carnitine in human proximal convoluted tubular (HPCT) cells in the kidney.

Plasma levels of octadecanoylcarnitine are decreased in patients with chronic fatigue syndrome, Alzheimer's disease and increased in patients with end-stage renal failure, as well as may be increased in congenital diseases such as celiac disease. Octadecanoylcarnitine is an inhibitor of glycine transporter 2 (GlyT2).

The product is used primarily as a control for MS/MS.



外观: 白色固體

分子量: 464.12

CAS 编号: 32350-57-5

分子式: $C_{25}H_{50}ClNO_4$

溶解度: DMF, DMSO, 乙醇

质量控制: NMR 1H 和HPLC-MS (95+%)

储存条件: 收到後 -20°C 避光保存 24 個月。運輸: 室溫最多可保存3週。乾燥。

法律声明: 本產品僅供研究目的提供和銷售。本產品並未經過食品、藥品、醫療器械、化妝品等領域的安全性和效力測試, 且未經明示或暗示授權用於其他任何用途, 包括但不限於體外診斷、人類或動物用途, 以及商業用途。